

Sensitivity analysis in economic simulations - a systematic approach

Claudia Hermeling
Tim Mennel*

Center for European Economic Research (ZEW)
P.O. Box 103443
D-68034 Mannheim, Germany

Preliminary version - do not quote!

April 2008

Abstract

Sensitivity analysis is an important part of quantitative modelling in economics and other empirical sciences. It studies how the variation in the output of a model (numerical or otherwise) can be apportioned, qualitatively or quantitatively, to different sources of variation. Thus, it serves to examine the robustness of numerical results with respect to input parameters, which is a prerequisite for deriving economic conclusions from them. In practice, modellers apply different methods, often chosen ad hoc, to do sensitivity analysis. This paper pursues a systematic approach. It formalizes deterministic and stochastic methods used for sensitivity analysis. Moreover, it presents the numerical algorithms to apply the methods, in particular, an improved version of a Gauss-Quadrature algorithm, applicable to one as well as multidimensional sensitivity analysis. The advantages and disadvantages of different methods and algorithms are discussed as well as their applicability.

*Corresponding Author, e-mail: mennel@zew.de. Financial Support by the EU project "Transust.Scan" is gratefully acknowledged.

1 Introduction

In economics as well as other model based sciences, a modeler has to do a sensitivity analysis to show the validity of results of his numerical model simulations. A sensitivity analysis is the study of how the variation in the output of a model (numerical or otherwise) can be apportioned, qualitatively or quantitatively, to different sources of variation in input parameters. It thus allows for an assessment of the robustness of numerical results, as it translates the range (confidence intervals) of fundamental (input) parameters into the model into ranges (confidence intervals) of economic (output) variables. The econometrician Edward Leamer makes it quite clear: "A fragile inference is not worth taking seriously. All scientific disciplines routinely subject their inferences to studies of fragility. Why should economics be different? ... What we need are organized sensitivity analyses." (Leamer, 1985)

In the context of CGE models, we ask whether the choice of basic parameters of the model, e.g. elasticities or time preference parameters, lead to stable equilibrium values of economic variables, e.g. GDP or labor participation. Usually, we refer to the equilibrium of the benchmark scenario. Quite importantly, a sensitivity analysis depends on the existence of equilibria for a sufficient range of parameters: If the model is not solvable for parameter values close to the ones we have chosen as benchmark values, model results are instable and thus worthless.

Basically, there are two methodological approaches to sensitivity analysis: a deterministic and a stochastic approach. Deterministic sensitivity analysis assumes that the tuple of basic parameters is an element of a given subset of all possible parameter choices. It seeks to determine upper and lower bounds on the corresponding subset of economic outcomes of the model. Stochastic sensitivity analysis treats the vector of parameters as a stochastic variable with a given distribution, rendering economic equilibria of the model into stochastic variables. It aims at calculating the first moments of these variables, with the variance indicating the robustness of the results. The two approaches are presented in section 2.

The choice a modeler has to make in a sensitivity analysis is, however, not only a methodological, but also a numerical one. Sensitivity analysis can involve more or less calculations of equilibria, so that usually there is a trade-off between accuracy and calculation time. This holds already true for a comparison of the deterministic and the stochastic approach, and is particularly relevant for the case of a multidimensional sensitivity analysis. In section 3, we present and discuss different algorithms. In particular, an improved version of an algorithm based on

Gauss-Quadrature is developed.

Section 4 presents a simple CGE model in Markusen's (2002) spirit. We conduct a sensitivity analysis with respect to demand elasticities and discuss the results obtained with different algorithms.

2 Theory of Sensitivity Analysis

2.1 Mathematical Preliminaries

Before conducting a sensitivity analysis, indeed even before implementing a numerical model, the modeller has to understand whether his model is in fact solvable. Proving existence (and sometimes uniqueness) of economic equilibria is a challenge of its own. There exist, however, a set of mathematical theorems that - if applicable - guarantee the existence of solutions. For example, the existence of an equilibrium in a general equilibrium framework is proved by recurring on Kakutani's fix-point theorem (cf. Mas-Collel et al. 1995, chapter 10). In the sequel, we assume that models are (uniquely) solvable for at least some parameter values and discuss under which circumstances a sensitivity analysis is possible.

An equilibrium of a computable general equilibrium (CGE) model takes the mathematical form of a solution to a system of (non-linear) equations

$$G(x^*, a) = 0,$$

where $x^* \in \mathbf{R}^n$ is a vector of (equilibrium) state variables of the economy (such as capital or wage) and $a \in \mathbf{R}^d$ a vector of parameters of the economy (such as demand elasticity or time preference). G is a continuously differentiable function

$$\mathbf{R}^n \times \mathbf{R}^d \rightarrow \mathbf{R},$$

that consists of first order conditions and (budget) constraints. This CGE model will be our standard example. Note, however, that the methods for sensitivity analysis presented can be generally be applied to economic models: The decisive distinction is the one between economic state variables x and basic parameters a .

Sensitivity analysis is concerned with the effect that (minor) changes of basic parameters Δa have on equilibrium state variables Δx^* . This notion will be formalized subsequently. At this point we ask under what circumstances an equilibrium x^{**} exists for a parameter value a' in a neighborhood of a , e.i. $a' \in$

$B_\epsilon(a) = \{\tilde{a} \text{ s.t. } |\tilde{a} - a| < \epsilon\}$. This is an important question: The existence of economic equilibria in a neighborhood of a is the theoretical prerequisite for sensitivity analysis. The implicit function theorem gives a definite answer.

Theorem 1 (Implicit function theorem) *If $\det|\nabla_{x^*}G(x^*, a)| \neq 0$, then there exists an open neighbourhood $\mathcal{U}(a) \subset \mathbf{R}^d$ of a and a continuously differentiable function $h : \mathcal{U}(a) \rightarrow \mathbf{R}^n$ that maps any vector of parameters on the corresponding equilibrium vector.*

Proof. Rudin (1976), theorem ? ■

Thus we learn that there is some $\epsilon > 0$ so that the existence of equilibria in a ϵ -neighborhood of a is guaranteed whenever G is a regular function at a . Generally speaking, we would expect G to be regular as long as first order conditions and constraints are independent. While it may be difficult to prove the assertion in some cases, it can be checked without problem numerically.

We have formulated the implicit function theorem for the case of CGE models. Similar formulations can be given for partial equilibrium models that are characterized by first and second order conditions and, possibly, additional constraints. In the case of economic optimization problems, the role of the implicit function theorem is taken by the theorem of the maximum (cf.).

So far we have neglected the notion of uniqueness of equilibria. While in principle sensitivity analysis can be conducted in the presence of multiple equilibria as well, uniqueness facilitates the analysis considerably. It is usually ensured by adequate convexity assumptions (cf. MasCollé et al. (1995) p.?). In the more general case of multiple equilibria, caution is warranted. In this case,

$$h : \mathcal{U}(a) \rightarrow \mathcal{P}(\mathbf{R}^n)$$

is a correspondence, mapping the vector of parameters into a set of solutions

$$h(a) = \{x_1^*, x_2^*, \dots, x_m^*\}.$$

Ignoring multiplicity can seriously blur a sensitivity analysis whenever a numerical solver 'jumps' from a solution x_j^* to some other solution x_i^* along changes of underlying parameters a . In that case, sensitivity of an equilibrium with respect to the basic parameters can be seriously exaggerated: Instead of following the initial equilibrium x_j^* along a continuous path for changes of a (cf. Judd, page), a numerical discontinuity occurs and the new equilibrium \tilde{x}_i^* is more distant to x_j^* than the correct equilibrium \tilde{x}_j^* . The best provision against such fallacies is the

calculation of all equilibria along the path of change; however, this can entail a considerable computational effort. At this point, we will not discuss the issue any further.

2.2 Deterministic Sensitivity Analysis

Sensitivity analysis is sometimes called robustness analysis. This term highlights its motivation: Assuming that we do not know the basic set of parameters exactly, how robust are the economic state variables in an equilibrium with respect to changes in the parameters? The **deterministic approach** to sensitivity analysis states is that there exists one true vector of economic parameters $a^* \in \mathbf{R}^d$, but that -instead of a^* - we only know its neighbourhood \mathcal{A} . Usually, we choose one vector of parameters $\hat{a} \in \mathcal{A}$ and call it the *benchmark scenario*. The point of sensitivity analysis then is to investigate whether equilibria vary considerably across $h(\mathcal{A})$ in comparison to the benchmark equilibrium $h(\hat{a})$.

Mathematically speaking, deterministic sensitivity analysis amounts to a geometric problem: Determine the relation of the volume of the image of \mathcal{A} under h and the size of $h(\hat{a})$, weighted with a scaling factor w_k in each dimension¹

$$\frac{\text{vol}(im(w * h))}{\|w * h(\hat{a})\|} = \frac{\int_{\mathcal{A}} \sqrt{\det(\partial_{ij}(w * h(a)))} da_1 \dots da_d}{(w * h(\hat{a}), w * h(\hat{a}))},$$

where the vector w specifies the relative weight we want to attach to the different economic variables in equilibrium². The findings of the model are robust whenever the relation is sufficiently small, where the assessment of sufficiency is left to the reader. The formal definition we have just given is a generalization of the more familiar notion of sensitivity analysis in one dimension: E.g. we might ask how big is maximal interval of values of GDP engendered by a model for a given interval of demand elasticities. We will discuss the issue in more detail in the next section where we present the *peacemeal approach* to sensitivity analysis.

Under some circumstances, we can say that the sensitivity of the model for a set of parameters \hat{a} is approximated by the relation of

¹By ' $*$ ' we denote component-wise multiplication of vectors.

²We do not have to attach a weight to an economic variable explicitly - instead we assess sensitivity by comparing the effect of changes in a on each dimension of the equilibrium state variable x . Implicitly, however, we will attach some weight to each dimension by accepting a certain outcome of the sensitivity analysis.

$$(w, \nabla h(\hat{a})) / (w, h(\hat{a})).$$

This Taylor-expansion view is used in linear programming (Flavell and Salkin, 1975). However, for non-linear programming it is usually impracticable and unreliable.

2.3 Stochastic Sensitivity Analysis

The **stochastic approach** to sensitivity analysis takes a different view of the basic problem: It treats the vector of basic parameters as a stochastic variable a with a given distribution $G(a)$ of $a \in \mathcal{A}$. While somewhat counterintuitive in the first place, the approach is in line with econometric estimations. These do not only produce mean values for parameters such as demand elasticities, but confidence intervals and higher moments for them. Under stochastic sensitivity analysis, h becomes a mapping onto a stochastic variable $x^* = h(a)$ of equilibria.

We then calculate the mean and the variation of the equilibrium vector x^* :

$$m = E[h(a)] = \int_{\mathcal{A}} h(a) dG,$$

$$v = Var[h(a)] = E[(h(a) - m)^2] = \int_{\mathcal{A}} (h(a) - m)^2 dG.$$

Attaching different weights to different economic variables, the stochastic sensitivity analysis assesses the size of

$$\sum_{k=1}^n w_k \frac{v_k}{m_k},$$

where index k is running over the dimension of x . In words: Given a distribution of basic parameters, we investigate the most likely equilibrium (the mean). We assess its robustness by assessing the relative size of the variance of equilibria with respect to the mean, possibly attaching different weights to different economic variables.

3 Practical Sensitivity Analysis

Having formalized the notion of sensitivity analysis in the preceding section we now present the practical implementation of the (somewhat abstract) concepts.

For the sake of clarity in the sequel we assume that all dimensions are weighted equally, thus dropping the vector w from all formulae.

3.1 The Peacemeal Approach

In a **peacemeal approach** to sensitivity analysis, we calculate

$$\Delta = \max_{a_i, a_j \in \{a_1, \dots, a_M\}} |h(a_i) - h(a_j)|$$

for a set of representative parameters $a_i \in \mathcal{A}$. The relation of Δ to the weighted benchmark equilibrium $h(\hat{a})$ is used to assess the sensitivity of the model at equilibrium $h(\hat{a})$. The peacemeal approach is kindred to a deterministic sensitivity analysis. But instead of calculating the volume of \mathcal{A} under h , it focusses on the maximal intervals of economic variables engendered by the set of parameters $a \in \mathcal{A}$.

If the set contains $\underline{a} = \mathbf{argmin}_{\mathbf{a} \in \mathcal{A}} \mathbf{h}(\mathbf{a})$ and $\bar{a} = \mathbf{argmax}_{\mathbf{a} \in \mathcal{A}} \mathbf{h}(\mathbf{a})$, then the following inequality holds (to facilitate the presentation and without loss of generality, we set $n = 1$):

$$\frac{\text{vol}(\text{im}(h))}{|h(\hat{a})|} \leq \text{vol}(\mathcal{A}) \frac{h(\bar{a}) - h(\underline{a})}{|h(\hat{a})|} = \text{vol}(\mathcal{A}) \frac{\Delta}{|h(\hat{a})|}.$$

A peacemeal approach can give a good idea of the sensitivity of the model if the set of parameters $a_i \in \mathcal{A}$ is sufficiently representative.

As an example, let both $n = 1$ and $d = 1$. Then

$$h : [\underline{a}, \bar{a}] \rightarrow [\underline{x}, \bar{x}]$$

for some scalar parameters \underline{a} , \bar{a} , \underline{x} and \bar{x} . If h is monotonously increasing, then

$$\frac{\text{vol}(\text{im}(h))}{h(\hat{a})} \leq (\bar{a} - \underline{a}) \frac{h(\bar{a}) - h(\underline{a})}{h(\hat{a})}.$$

In this specific case, the peacemeal approach and the more formal definition of deterministic sensitivity analysis given in the first section virtually coincide. Generally speaking, while our formal definition captures the intention of sensitivity analysis more accurately, the peacemeal approach is more practicable.

3.2 The Monte-Carlo Approach

The **Monte-Carlo** approach is the first of two practical implementations of stochastic sensitivity analysis. Both mean and variance of equilibrium x^* are approximated in the following way: Draw a (large) set of realisations $\{a_1, \dots, a_M\}$ from the distribution $G(\mathcal{A})$ and calculate

$$m = E[h(a)] \approx \frac{1}{M} \sum_{i=1}^M h(a_i) = \tilde{m},$$
$$v = Var[h(a)] \approx \frac{1}{M} \sum_{i=1}^M (h(a_i) - \tilde{m})^2 = \tilde{v}.$$

The sums of the right-hand side converge stochastically to the true values of m and v . Beyond mean and variance of the stochastic variable x^* , we can easily approximate its distribution $h \circ G$. A great disadvantage of the Monte-Carlo approach is that in order to assure convergence, the number of draws M has to be high and thus the approximation is numerically costly. This is a problem in particular when the space of parameters a is high dimensional - the curse of dimensionality drives up the number of necessary draws exponentially (cf. Judd).

3.3 The Gauss-Quadrature Approach

The second way of practically implementing stochastic sensitivity analysis is by **Gauss quadrature** - in fact a numerical method to approximate integrals (cf. Stoer). Remember that we intent to approximate mean and variance, that are defined by integrals of the distribution of basic parameters a . We want to do so using a rather small number L of function evaluations $h(\cdot)$.

Essentially, the Gauss quadrature gives us nodes x_i and weights ω_i to approximate the (one dimensional) integral

$$\int_a^b f(x)\omega(x)dx \approx \sum_{i=1}^L \omega_i f(x_i). \quad (1)$$

In our specific case, we look for nodes a_i and weights g_i to approximate mean and variance of equilibria:

$$m = \int_{\mathcal{A}} h(a)dG \approx \sum_{i=1}^L g_i h(a_i) = \tilde{m},$$

$$v = \int_{\mathcal{A}} (h(a) - m)^2 dG \approx \sum_{i=1}^L g_i (h(a_i) - \tilde{m})^2 = \tilde{v},$$

where again, the dimension of economic variables is set to $n = 1$ (we present the generalisation to several variables below).

In the following, we develop a version of Gauss quadrature new to computational economics, in that it builds on *orthogonal polynomials*. While somewhat complicating the straightforward Gauss quadrature algorithm commonly used in economics (cf. Arndt 1996) conceptually, our approach simplifies the computation of a sensitivity analysis in cases of standard probability distributions, increasing the approximation quality at the same time. This is possible because the optimal nodes x_i turn out to be zeros of orthogonal polynomials. They have to be linearly transformed to fit the respective interval but can otherwise be taken from an existing table. In contrast, in Arndt's (1996) algorithm, the nodes are the solution of a system of non-linear equations.

To define orthogonality in this context, let the distribution $G(a)$ be represented by a weight function $g(a)$. Then the expression

$$(f_1, f_2)_g = \int_{\mathcal{A}} f_1(a) f_2(a) g(a) da$$

defines a scalar product $(\cdot, \cdot)_g$. We refer to orthogonality with respect to this scalar product. The following lemma holds:

Lemma 2 (Gram-Schmidt, Weierstrass) *For any scalar product (\cdot, \cdot) on the space of continuous functions $\mathcal{C}([a, \bar{a}])$, there is a complete system of orthogonal polynomials $\{p_0, p_1, \dots | (p_i, p_j) = 0, i \neq j\}$.*

Proof.

For any given scalar product, orthogonal polynomials can be constructed from monomials $1, x, x^2, \dots$ by the Gram-Schmidt procedure

$$p_0 \equiv 1 \quad p_i(x) = x^i - \sum_{j=1}^{i-1} \frac{(p_j, x^i)}{(p_j, p_j)} p_j.$$

We thus obtain an infinite sequence of orthogonal polynomials. As for completeness, we know that the polynomials $(p_0(x), p_1(x), \dots, p_n(x))$ span the same linear subspace of the space of continuous functions as the monomials $(1, x, x^2, \dots, x^n)$.

Consequently, we can apply Weierstrass' approximation theorem which states that the space of polynomials is dense in the space of continuous functions (cf. Rudin 1976, ch. 2) and the completeness of the family of orthogonal polynomials ensues.

■

There are well known examples of orthogonal polynomials, the best known being Legendre, Tchebychev, Laguerre and Hermite polynomials.

Examples of families of orthogonal polynomials

Name	$g(x)$	$[a, b]$	Definition
Legendre	1	$[-1, 1]$	$P_k(x) = \frac{(-1)^k}{2^k k!} \frac{d^k}{dx^k} [(1-x^2)^k]$
Tschebyscheff	$(1-x^2)^{-\frac{1}{2}}$	$[-1, 1]$	$T_k(x) = \cos(k \cos^{-1}(x))$
Laguerre	$\exp(-x)$	$[0, \infty)$	$L_k(x) = \frac{\exp(x)}{k!} \frac{d^k}{dx^k} (x^k \exp(-x))$
Hermite	$\exp(-x^2)$	$(-\infty, \infty)$	$H_k(x) = (-1)^k \exp(x^2) \frac{d^k}{dx^k} (\exp(-x^2))$

The proof of lemma is constructive, so that for any weight function $g(a)$ orthogonal polynomials can be constructed from monomials $1, x, x^2, \dots$. For a general distribution probability distribution $G(a)$, their calculation can entail considerable

To proceed, we need one property of orthogonal polynomials.

Lemma 3 *The zeros $\{a_1, a_2, \dots, a_l\}$ of $p_l(a)$ are real and distinct.*

Proof. Stoer ■

It is because that they are real and distinct that the zeros of a orthogonal polynomial are a possible choice of nodes for the evaluation of the approximation formula 1. The following theorem shows that they are indeed a good choice.

Theorem 4 (Stoer) *Let $\{a_1, a_2, \dots, a_l\}$ be the zeros of $p_l(a)$ and g_1, \dots, g_l be the solution of the system of linear equations*

$$\sum_{i=1}^n g_i p_k(a_i) = \begin{cases} (p_0, p_0) & : k = 0 \\ 0 & : k = 1, 2, \dots, l-1 \end{cases}$$

Then $g_i > 0$ for $i = 1, 2, \dots, l$ and

$$\int_{\underline{a}}^{\bar{a}} p(a)g(a)da = \sum_{i=1}^l g_i p(a_i)$$

for all $p \in \Pi_{2l-1} = \ll p_0, \dots, p_{2l-1} \gg$.

In words: For a given weight function $g(a)$ (i.e. probability distribution G), we calculate the zeros a_1, \dots, a_l of the corresponding orthogonal polynomial of degree l . Calculating the weights g_1, \dots, g_l from a suitable system of linear equations, we obtain an integration formula of type 1 that integrates polynomials up to degree $2l-1$ exactly.

Thus, for our purpose of numerical integration, we have to calculate the zeros of orthogonal polynomials and weights corresponding to the probability distribution G with weight function $g(a)$. *However, we have to do so only once for a given G .* While in general the numerical determination of the zeros of orthogonal polynomials for a given distribution may be tricky, in the case of standard probability distributions we have no problem. A look at the table of orthogonal polynomials confirms that for uniform distributions, we can use Legendre polynomials, and Hermite polynomials for normal distributions. This facilitates our task considerably: We can either (easily) calculate the zeros numerically from the defining formulae for Legendre or Hermite polynomials, or take these from published tables. The next section shows an application of the method presented here to a simple example.

In higher dimensions $n > 1$, integrals can be approximated by product rules, combining one-dimensional nodes and weights:

$$\begin{aligned} & \int_{\underline{a}^1}^{\bar{a}^1} \dots \int_{\underline{a}^d}^{\bar{a}^d} f(a^1, \dots, a^d) g^1(a^1) \dots g^d(a^d) da^d \dots da^1 \\ & \approx \sum_{i_1=1}^n \dots \sum_{i_d=1}^n g_{i_1}^1 \dots g_{i_d}^d f(a_{i_1}^1, \dots, a_{i_d}^d) \end{aligned}$$

In a later version of this paper, we will provide the formulae for a higher-dimensional application of our Gauss-Quadrature approach. It is essentially straightforward, applying the one-dimensional procedure step by step for each dimension. It is in dimensions higher than one - when a joint distribution of basic parameters is inserted into the sensitivity analysis, that Gauss-Quadrature integration

has a great advantage over Monte-Carlo simulations, as the evaluation of nodes increases exponentially with each dimension, making MC simulations simply to expensive.

4 A Simple Example

We consider a two-by-two closed economy in the spirit of Markusen (2002). It represents an economy with two commodity goods, X and Y, two factors (capital K and labor L), and one single representative agent. The goods are produced through constant returns to scale production activities which combine primary factor inputs. We use a balanced equilibrium data set given by the square accounting matrix below. The accounts labelled X and Y in this matrix refer to markets for final commodities. Account W correspond to final consumption. The RA account corresponds to the representative agent. It defines both the endowment and expenditures for the models's single representative agent.

Accounting matrix with benchmark flows:

Markets	X	Y	W	RA	Row sum
PX	100		-100		0
PY		100	-100		0
PW			200	-200	0
PL	-40	-60		100	0
PK	-60	-40		100	0
Column sum	0	0	0	0	

The accounts of the matrix do not by themselves completely characterize a general equilibrium framework because they provide a variety of benchmark value shares. A model formulation additionally relies on assumptions about elasticities of substitution in the various sectors. These are the parameters with respect to which we want to conduct a sensitivity analysis.

In our model we have three elasticities of substitution for the three production activities: $esubx$ denotes the elasticity of substitution between inputs to X production, $esuby$ the elasticity of substitution between inputs to Y production and $esubw$ the one between inputs to final consumption.

We choose $esubx = esuby = esubw = 0.5$ and use a GAMS program to calibrate the model as replication of the benchmark equilibrium. Then we introduce an exogenous labor tax on good X by setting $TAX_LX = 1$.

In the following we want to compare two different approaches for the sensitivity analysis: the Monte-Carlo (MC) analysis and the Gauss-Quadrature (GQ). We assume that the elasticity of substitution e_{subx} is uniformly distributed on the interval 0.25 and 0.75 and that the other elasticities are constant and equal to 0.5. Due to the assumption of uniform distribution we choose Legendre polynomials for the Gauss-Quadrature.

For the sensitivity analysis with the Monte-Carlo approach we randomly draw values for e_{subx} from the interval [0.25,0.75]:

$$e_{subx} = \text{UNIFORM}[0.25, 0.75].$$

We calculate the corresponding equilibrium values for X, Y and W: X.L, Y.L and W.L. Given the solution from the benchmark scenario with x_{bench} , y_{bench} and w_{bench} the new results are normalized:

$$\text{result}_x = \text{ROUND}(100 * (X.L - x_{bench})/x_{bench}).$$

For y and w, this is done accordingly. By summing up

$$\text{results}(i) = (\text{result}_x(i), \text{result}_y(i), \text{result}_w(i))$$

over all drawings i and dividing by the number of draws we calculate the mean and hereupon the variance of the economic variables.

For the sensitivity analysis using Gaussian Quadrature with Legendre polynomials we use a MATLAB routine to calculate the zeros of Legendre polynomials `legendrenodes` and the weights `legendreweights`. Hereby, we use the recursive formula for polynomials given by the Gram-Schmidt procedure, calculate the zeros for these polynomials and solve the equation system given in Theorem 4 with respect to the corresponding weights. We save the zeros and weights in a `gdx` file. Alternatively, schedules with zeros and weights for Legendre polynomials can also be found in the internet or in math books. Once this `gdx` file is created, it can be used to run sensitivity analysis for every CGE model where the parameters are assumed to be uniformly distributed.

In our GAMS program, the variables `legendrenodes` and `legendreweights` are loaded and transformed into the variables `grid` and `weights`. For the program we choose `maxdegree` which is the degree of the maximal Legendre polynomial that we want to consider. It also refers to the rows with zeros and weights that are loaded from the `gdx` file. A higher `maxdegree` raises the accuracy of the Legendre approach but is computationally more expensive.

We have to transform `legendrenodes` linearly from interval $[-1, 1]$ to `grid` on interval $[a, b] = [0.25, 0.75]$ because the standard Legendre polynomials are defined on the interval $[-1, 1]$: $\text{grid}(i) = (b - a) * (\text{zerosval}('maxdegree', i) + 1)/2 + a$. `zerosval('maxdegree', i)` denotes the i -th zero point for a Legendre polynomial of degree `maxdegree` which is loaded from `legendrenodes`. Now we calculate the equilibrium values for X , Y and W by a loop of the model, varying the elasticity `esubx` across `grid`. We calculate again the means and variances of economic variables by summing up results, weighted by `weights` which can be drawn from the `gdx` file directly:

$$\text{mean}(\text{result}_x) = \text{SUM}(i, \text{weights}(i) * \text{result}_x(i)),$$

$$\text{variance}(\text{result}_x) = \text{SUM}(i, \text{weights}(i) * \text{sqr}(\text{result}_x(i))) - \text{sqr}(\text{mean}(\text{result}_x)).$$

For y and w , this is done accordingly.

Sensitivity w.r.t. `esubx`: Mean

Name (Runs)	MC mean (100)	MC mean (500)	MC mean (2000)	GQ mean (10)	GQ mean (20)	GQ mean (40)
X	-8.990	-9.010	-9.023	-9.036	-9.026	-9.030
Y	7.582	7.578	7.575	7.576	7.568	7.570
W	-1.396	-1.410	-1.417	-1.426	-1.427	-1.421

Sensitivity w.r.t. `esubx`: Variance

Name (Runs)	MC var. (100)	MC var (500)	MC var (2000)	GQ var (10)	GQ var (20)	GQ var (40)
X	0.027	0.031	0.031	0.031	0.031	0.031
Y	0.004	0.004	0.005	0.004	0.004	0.005
W	0.014	0.015	0.016	0.018	0.015	0.016

The matrix above presents the result for our model. The number of model evaluations which is proportional to the computation time are given as following: $M_{MC} = 100$, $M_{MC} = 500$ and $M_{MC} = 2000$ for the Monte Carlo approach and $l_{GQ} = 10$, $l_{GQ} = 20$ and $l_{GQ} = 40$ for the Gauss-Quadrature. It can be seen directly from the matrix that the Gauss-Quadrature considerably reduces the computation time.

The sensitivity analysis can be done in a similar way for the parameters Y and W separately or for all parameters simultaneously. In the later case, the elasticities of substitution are drawn randomly but independently from given intervals $[a_x, b_x]$, $[a_y, b_y]$ and $[a_w, b_w]$ when using the Monte Carlo approach. For the Gauss-Quadrature three grids are loaded and transformed from the gdx file. The equilibrium values are calculated by three loops over the three grids. Mean and variance can be derived by weighting the results with the associated weights:

$$\text{mean}(\text{result}_x) = \text{SUM}(i, \text{SUM}(ii, (\text{SUM}(iii, \text{weights}(i) * \text{weights}(ii) * \text{weights}(iii) * \text{result}_x(i, ii, iii))))))$$

and

$$\text{variance}(\text{result}_x) = \text{SUM}(i, \text{SUM}(ii, \text{SUM}(iii, \text{weights}(i) * \text{weights}(ii) * \text{weights}(iii) * \text{sqr}(\text{result}_x(i, ii, iii)))))) - \text{sqr}(\text{mean}(\text{result}_x))$$

5 Conclusion

Due to its general importance in economic modelling, sensitivity analysis merits a systematic understanding by economic modellers. This paper hopes to contribute to such an understanding and to serve as a guide in applying the appropriate algorithm. Dimensionality of sensitivity analysis (the number of parameters that are varied simultaneously) makes an important difference. Moreover, it is demonstrated that generally stochastic sensitivity analysis gives more and better insights than deterministic sensitivity analysis. However, it is also more burdensome computationally. Comparing stochastic methods, the paper shows that Monte-Carlo methods are easily applicable, but computationally expensive. Gauss-Quadrature methods reduce the computational burden and thus are suitable for higher dimensional problems.

References

- [1] Arndt, C. (1996), "An Introduction to Systematic Sensitivity Analysis via Gaussian Quadrature", GTAP Technical Paper No. 2, Purdue University
- [2] Flavell, R. and G. R. Salkin (1975), "An Approach to Sensitivity Analysis", *Operations Research Quarterly*, Vol. 26 (4), pp. 857-866
- [3] Judd, K. (1998), "Numerical Methods in Economics", MIT Press
- [4] Leamer, E. (1985), "Sensitivity Analyses would Help", *American Economic Review*, Vol. 75 (3), pp. 308-313
- [5] Mas-Colell, A., Whinston, M. D. and J. R. Green (1995), "Microeconomic Theory", Oxford University Press
- [6] Markusen, J. A. (2002) "General-Equilibrium Modeling using GAMS and MPS/GE", mimeo, University of Colorado
- [7] Rudin, W. (1976), "Analysis", McGraw-Hill
- [8] Stoer, J. and R. Bulirsch (1990) "*Introduction to Numerical Analysis*", Springer